



ULTRACARE®

Oral Anesthetic Gel

Description:

Ultracare is a 20% w/v (17.9%) benzocaine oral anesthetic gel preparation in a water-soluble glycol base. Ultracare is designed for rapid onset (10-30 seconds). Anesthesia usually lasts 8-10 minutes. Ultracare is NOT for injection.

Uses:

For the temporary relief of occasional minor irritation and pain, associated with:

- minor dental procedure, including local anesthesia injections, scaling or prophylaxis. It may also be used to temporarily relieve discomfort of the oral mucosa associated with making impressions, intraoral radiographs, placing a rubber dam clamp or other manipulation of the oral mucosa.
- sore mouth and throat
- minor injury of the gums
- canker sores
- minor irritation of the mouth or gums caused by dentures or orthodontic appliances

MAXIMUM DOSAGE: 0.4g (0.4ml) per day

Procedure:

- I.** Preloaded 1.2ml syringes:
 - a.** Remove luer lock cap from syringe.
 - b.** Attach a Black Micro® 20-gauge Tip or Turquoise Capillary Tip securely onto syringe.
 - c.** Slowly express gel to the target tissue.
 - d.** For anesthetizing periodontal pockets:
 - i.** Slowly apply a small amount to one quadrant at a time to the gingival margin or directly into the periodontal pockets and proceed with sub-gingival scaling.
-

2. IndiSpense delivery:
 - a. Remove cap from IndiSpense syringe.
 - b. Express desired amount of anesthetic to cotton applicator (Fig. 2) or into a dappen dish. Use a clean applicator with each area application. Re-cap the IndiSpense syringe after each use.
 - c. Using a dappen dish allows for re-application with the same applicator without the risk of cross-contamination. Re-cap the IndiSpense syringe after each use.
3. IndiSpense delivery to empty 1.2ml syringe:
 - a. Attach a 1.2ml syringe to the end of IndiSpense by turning the luer lock of the 1.2ml syringe snugly onto male thread of Indispense syringe.
 - b. With palm, press plunger of IndiSpense syringe gently while guiding the 1.2ml plunger outward to desired fill.
 - c. RE-cap the IndiSpense syringe after each use.
 - d. See instruction for 1.2ml syringe use.

Directions:

Adults and children 2 years of age and older:

- Apply (a thin film) to affected area and allow to remain in place approximately 1 minute.
- Rinse and expectorate to clear the oral cavity of residual product.

For topical anesthesia prior to injection:

- Apply a thin film to injection site(s) allowing the gelled cotton applicator to remain in place at least 1 minute.
- Rinse and suction the oral cavity and continue with administering local anesthetic or with dental procedure.

For anesthetizing periodontal pockets:

- See instruction for 1.2ml use.




Fig. 1 Hold plunger in palm of hand for optimum control.



Fig. 2 Deliver Ultracare from the large, no-waste IndiSpense syringe to cotton applicator, cotton roll, etc., for placement.

Warnings and Precautions:

1. For Professional use only.
 2. For external use only.
 3. Keep out of reach of children.
 4. Ultracare is NOT for injection.
 5. Do not use on patients:
 - a. With a known sensitivity to PABA, sunscreen or to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.
 - b. With methemoglobinemia or those taking sulfonamides.
 - c. Do not use on children under 2 years of age.
 6. Pregnancy and Lactation: Safety is not known in pregnancy/nursing women.
 7. Avoid contact with eyes.
 8. If applying intraorally from a syringe, verify flow on a gauze pad or mixing pad prior to applying. If resistance is met, replace tip and re-check.
 9. If an excessive amount of gel is swallowed, get medical help or contact a Poison Control Center immediately.
 10. Do not exceed maximum recommended dosage.
 11. Phenylketonurics: Contains Phenylalanine—3mg per gram
 12. See state or country guidelines for proper disposal of empty jar, syringes, applicator and tips.
 13. To avoid cross-contamination of the prefilled syringes, use a disposable syringe cover; re-cap, and wipe syringe with an intermediate disinfectant between uses. If these measures are not taken, the syringes should be considered single-use. Tips are disposable. To avoid cross-contamination, do not re-use tips.
-

For immediate reorder and/or complete descriptions of Ultradent's product line, refer to Ultradent's catalog or call Toll Free:
1-800-552-5512.
Outside U.S. call (801) 572-4200.

Ultracare



Hazard Rating
4 = Severe
3 = Serious
2 = Moderate
1 = Slight
0 = Minimal

For product SDS please see our website: www.ultradent.com

All Ultradent syringes have an expiration date stamped on the side of the syringe consisting of one letter and three numbers. The letter is a lot number used for manufacturing purposes and the three numbers are the expiration date. The first two numbers are the month, and the third number is the last number of the year.

Key:




Use by Date



See Instructions
For Use



Keep out of
reach of children



Store at room
temperature



Health
Hazard



Do Not Reuse
to Avoid Cross
Contamination

REF

Catalog Number

LOT

Batch Code

© Copyright 2017 Ultradent Products, Inc. All Rights Reserved.

U2 ULTRADENT
PRODUCTS, INC.

Manufactured by:

Ultradent Products Inc.

 505 West Ultradent Drive (10200 South), South Jordan, Utah 84095, USA
Made in USA

#10057.15 052417